

Cell Sorting

How it works?


You bring us sort sample(s) and appropriate controls and we sort the desired population within your sample.

Astrios: 5-lasers 355,405,488,561,630 nm; sort into tubes or plates; sort microparticles.

Remember, book your sort and submit a sort-form at least 2 days before sort-day

<https://dfcf.acs.analytical.unsw.edu.au/>

Flow chart



Sort buffers

Basic	Sticky cell	Low viability
1x PBS 2 mM EDTA up to 2% serum	1x Hanks BSS 5 mM EDTA 25 mM HEPES, pH 7 up to 2% serum	1x Hanks BSS 2 mM EDTA 25 mM HEPES, pH 7 10U DNase up to 2% serum

Next page, gotchas

Gotchas

There are a number of critical steps for successful cell sorts

- **Cells MUST be filtered**
 - Even if your cells don't need it.
 - **Consequence:** Machine clogs, you take a big hit to your yield. Lose sort time for the rest of your samples.
- **Bring correct controls**
 - **Action required:** Common controls: unstained, compensation, FMO
 - **Consequence:** We will have to guess which population to sort if we can even proceed with the sort.
- **Over confluent cells**
 - **Action required:** Keep your cells <70% confluent (adhesive), 8×10^5 / mL (suspension)
 - **Consequence:** single-cell suspension will form aggregates faster, causing machine blockages even if filtered. Cells will be starving and not behave as expected (unless you always starve your cells?).
- **Low viability samples**
 - Action required: Add DNase to sample
 - Consequence: DNA released by dead cells causes clumping
- **Low yield**
 - **Main Causes:**
 - Population of interest was not as represented as expected
 - Total cell count was lower than expected
 - Clumped cells – which those events were aborted
 - Machine capability (up to 40% loss (aria), 25% loss (astrios). This includes cells exploding on impact in the tube
 - **Solutions:**
 - Low % of desired cells; optimise transfection – don't just follow someone else's protocol, establish the best conditions for your work.
 - Cell loss; try to identify cause of loss (during washes, column), or increase input amount if possible
 - Clumped cells; avoid over-confluence, we can filter the sample just before the sort to reduce cells re-adhering, use DNase to break up low-viability samples
 - Exploding cells; Astrios delivers higher yields for plate sorting

Sort time; as predictable as pachinko.

The sweet spot for a reasonable sort time is to sort a population with greater than 1% sample representation.	0.1	1.0	10.0	30.0	% cells of interest in sample
	2,000				event rate / sec
	1666.7	166.7	16.7	5.6	minutes for 200k
	5,000				
	666.7	66.7	6.7	2.2	
	9,000				
370.4	37.0	3.7	1.2		

- Well, sort at 9000/s *all* the time!

- This is only possible if your cells are pristine. We usually run at 5000 events / s.